

GABAergic Modulation of Yawning Behavior

ENRIQUE DOGER, RUTH URBÁ-HOLMGREN, JOSÉ RAMÓN EGUIBAR
AND BJÖRN HOLMGREN

*Departamento de Ciencias Fisiológicas, Instituto de Ciencias
Universidad Autónoma de Puebla, Apartado Postal 406, Puebla, Pue., Mexico*

Received 27 February 1989

DOGER, E., R. URBÁ-HOLMGREN, J. R. EGUIBAR AND B. HOLMGREN. *GABAergic modulation of yawning behavior*. PHARMACOL BIOCHEM BEHAV 34(2) 237-240, 1989.—The hypothetical modulation by GABAergic neurons of yawning behavior in the rat was explored with GABA-active drugs. Gamma-acetylenic-GABA, a specific inhibitor of GABA-T, increases yawning frequency when injected at a dose of 7 mg/kg. Baclofen, a GABA_B agonist (3 mg/kg), inhibits yawning completely; GABA antagonists, bicuculline and picrotoxin, at subconvulsant doses, also decrease yawning. All drugs were injected intraperitoneally with the exception of apomorphine, which was injected subcutaneously. It is suggested that GABA_B receptors play a role in yawning behavior by modulating ACh release, and that GABA_A receptors may modify yawning frequency by modulating inhibitory influences on ACh neurons.

Yawning GABA GABA_A receptors GABA_B receptors

YAWNING is a discrete innate motor pattern widely represented in the behavioral repertoire among vertebrates (5, 17, 18). It occurs spontaneously at variable frequencies (2,4), and is subject to a complex set of neurotransmitter and hormonal influences (6, 12, 13, 24, 27-29, 35, 43). Yawning can be induced by cholinomimetic drugs such as physostigmine and pilocarpine, and is inhibited by scopolamine (35, 39, 40). It may also be evoked by low doses of apomorphine (APO), and other dopamine (DA) receptor agonists (-3PPP and bromocriptine) (20, 29, 32, 37, 38, 43). With higher doses of APO, which also act upon postsynaptic receptors, spontaneous and physostigmine-induced yawning are inhibited (20, 32, 37, 38). In spite of the fact that other substances, like the peptide hormones ACTH (12,13), α -MSH (12), prolactin (24) and oxytocin (3), are yawning inducers, several authors agree in assigning a crucial role in the regulation of yawning to the DA-acetylcholine (ACh) interaction (15, 37, 42).

Links between ACh and DA neurons have been properly described in the septo-hippocampal pathway and the striatum (1, 16, 34). It has been described that stimulation and lesion of these structures affect yawning frequency (10, 21, 25, 33, 43).

Gamma-aminobutyric acid (GABA) is the main inhibitory neurotransmitter in the principal efferent systems related to the nigro-striatal and the ventral tegmental-nucleus accumbens dopaminergic systems (26,31). Also, it is well known that GABAergic neurons regulate dopaminergic and cholinergic activities in these structures (30, 31, 41). Therefore, it seemed important to explore the role of the GABA system in the regulation of yawning behavior. Two different types of GABA receptors have been described, GABA_A and GABA_B. The former are sensitive to muscimol and THIP, and are antagonized by bicuculline. Receptors of this kind are mostly situated postsynaptically, and are coupled to chloride channels. The other type of GABA receptors, GABA_B, are not sensitive to bicuculline, their most specific agonist is baclofen; they are mostly presynaptic and are involved

in the modulation of ACh release (8, 11, 22, 23).

The experiments here reported have been performed with GABA-mimetic drugs and GABA antagonists. The use of two sublines of rats with high and low spontaneous yawning frequency has allowed us to study the effect of these GABA-active drugs, both on spontaneous and drug-induced yawning.

METHOD

Animals

Male Sprague-Dawley rats, 2-3 months old, 200-300 g in weight, from two sublines selectively bred to establish high (HY) and low (LY) yawning frequency were used. They were bred in our animal house under standard conditions: light-dark cycle 12:12 hr, lights on from 0700 to 1900; food (Purina chow) and water were supplied ad lib. The animals were weaned at the age of 30 days and maintained in groups of four in collective Plexiglas cages.

Drugs

The drugs used in the experiments were the following: gamma-acetylenic-GABA (GAG) (Merrel Labs, France); bicuculline methiodide (Sigma, USA); picrotoxin (Sigma, USA); tetrahydroisoxipiridonol HCl (THIP) (Research Biochem. Inc., USA); α -p-chlorophenyl-GABA (baclofen) (Ciba Geigy, Basel, Switzerland); physostigmine sulphate (Sigma, USA); apomorphine HCl (APO) (Chimimport, Bulgaria); and 3-(3-hydroxyphenyl)-N-n-propyl piperidine HCl (-3PPP) (Astra, Sweden). Bicuculline was dissolved in 0.1 N HCl and diluted with saline. The other drugs were directly dissolved in saline. Solutions were freshly prepared before the experiments. A standard injection volume of 0.2 ml/100 g body weight was used. APO was injected subcutaneously (SC). Other drugs were administered intraperitoneally (IP). Doses are expressed as mg (base)/kg body weight. GAG was injected eight




FIG. 1. Effect of GAG on spontaneous yawning frequency. Ordinate: number of yawns above basal rate per hour; mean basal yawning rate (MBYR) = 19.4 yawns/hr. Abscissa: GAG doses. $n = 10$ HY male rats per dose (Kruskal-Wallis test, $p < 0.05$, Mann-Whitney U-test, $*p < 0.05$). Vertical bars indicate standard error (SE).

hours before the observation period, -3PPP 15 minutes before, and all the other drugs immediately before yawn monitoring began. Animals were used only once. Statistical procedures will be mentioned with the results.

Behavioral Observations

Observations were performed with each animal placed in a transparent glass cylinder (diameter 19 cm, height 10 cm), the floor of which was covered with a sheet of clean filter paper, and the top with a Plexiglas plate, leaving a segment 1 cm wide open for ventilation. The observation periods were between 1700–1800 hr. Yawns were monitored by two observers sitting on opposite sides of the table on which the animals were placed. Usually eight rats were observed simultaneously.

RESULTS

GAG, a specific inhibitor of GABA-transaminase (the catabolic enzyme of GABA), exerts GABA-mimetic activity with a long latency. Its maximal effects on yawning are observed 8 hours after administration (9). Therefore, in these experiments, the rats were injected 8 hr before the behavioral observations.

The dose-response curve of GAG on spontaneous yawning was plotted with four doses (Fig. 1). A significant increase in yawning was obtained only with 7 mg/kg; with higher doses (15 and 25 mg/kg) the animals appeared very sleepy and hypodynamic. In order to study the effects of the drug on pharmacologically-induced yawning we used the following experimental design: 10 HY male rats, pretreated with GAG (7 mg/kg, 8 hr before), were injected with the yawning inducer and observed during 1 hr. Figure 2 shows the results when physostigmine (0.15 mg/kg IP), APO (0.05 mg/kg, SC) and -3PPP (10 mg/kg IP, 15 min before observation) were used as inducers. The increase in yawning frequency, observed in GAG-physostigmine-injected rats, was the only significant result obtained ($p < 0.05$, Mann-Whitney U-test).

In other experiments, we tested THIP, a GABA_A agonist, and baclofen, a GABA_B agonist, in both HY and LY rat sublines. Systemic administration of THIP, at 0.33, 1.0, 3.0 and 6.0 mg/kg, does not produce any change in yawning frequency (Kruskal-Wallis, $p > 0.05$). On the other hand, baclofen produces a clear inhibition of spontaneous yawning behavior (Fig. 3) (Kruskal-Wallis, $p < 0.05$; Mann-Whitney U-test, $p < 0.05$ or less). Baclofen also inhibits physostigmine-induced yawning (Fig. 3).

The effects of GABA antagonists, bicuculline and picrotoxin, were also tested in HY male rats. Both drugs, at subconvulsant doses, reduce spontaneous yawning frequency. With bicuculline the highest effect was obtained with 3 mg/kg, while 1 mg/kg




FIG. 2. Effect of GAG on pharmacologically-induced yawning. Ordinate: number of yawns above basal rate per hour; PHY = physostigmine, 0.15 mg/kg, IP, MBYR = 25.4 yawns/hr; -3PPP, 10 mg/kg, IP, MBYR = 19.3 yawns/hr; APO = apomorphine 0.05 mg/kg, SC MBYR = 22.1 yawns/hr; GAG 7 mg/kg, IP. $n = 10$ HY male rats per group (Mann-Whitney U-test, two-tailed, $*p < 0.05$). PHY and -3PPP results are referred to the left ordinate. For each combination of drugs, two groups of animals were compared: one of them was pretreated with GAG injected 8 hr before, the other with saline, and both received the indicated yawning-inducing drug. The three yawning-inducing drugs increase basal yawning rates significantly (Mann-Whitney U-test, $p < 0.05$ or less). Vertical bars indicate SE.

picrotoxin suppressed yawning completely (Fig. 4) (Kruskal-Wallis, $p < 0.05$; Mann-Whitney U-test, $p < 0.05$ or less).

DISCUSSION

In order to evaluate the possible role of the GABA system in the regulation of yawning we studied the effects of different GABA-active drugs upon this behavior.

The decrease in spontaneous yawning frequency observed with baclofen (GABA_B agonist) suggests that GABA neurons play a role in yawning regulation. This decrease may be caused by a lower central cholinergic tone, due to a presynaptic GABAergic modulatory action on cholinergic terminals. This sort of modula-




FIG. 3. Effect of baclofen on spontaneous and physostigmine-induced yawning. Ordinate: number of yawns observed during 1 hr. Abscissa: baclofen doses (mg/kg) IP. PHY = physostigmine 0.15 mg/kg, IP, $n = 12$ male rats per dose. LY and HY dose-response curves: Kruskal-Wallis test, $p < 0.05$; Mann-Whitney U-test, * and $**p < 0.05$, and $p < 0.02$ when compared with the control. Vertical bars indicate SE.

- Pharmacol. Exp. Ther. 240:294–297; 1987.
8. Bowery, G. Baclofen: 10 years on. In: Lambie, J. W.; Abbott, A. C., eds. Receptors again. Amsterdam, The Netherlands: Elsevier Science Publishers; 1984:195–202.
 9. Doger, E.; Urbá-Holmgren, R.; Holmgren, B. Effect of gabaergic drugs on the yawning behavior of the rat. *Soc. Neurosci. Abstr.* 11:1289; 1985.
 10. Dourish, C. T.; Hutson, P. H. Bilateral lesions of the striatum induced with 6-hydroxydopamine abolish apomorphine-induced yawning in rats. *Neuropharmacology* 11:1051–1055; 1985.
 11. Enna, S. J.; Karbon, E. W. In: Olsen, R. W., Venter, J. C., eds. GABA receptors: An overview in benzodiazepines/GABA receptors and chloride channels: Structural and functional properties. New York: Alan R. Liss, Inc.; 1986:41–56.
 12. Ferrari, W. Behavioral changes in animals after intracisternal injection with adrenocorticotrophic hormone and melanocyte stimulating hormone. *Nature* 181:925–926; 1958.
 13. Gessa, G. L.; Pisano, M.; Vargiu, L.; Crabai, F.; Ferrari, W. Stretching and yawning movements after intracerebral injection of ACTH. *Rev. Can. Biol.* 26:229–236; 1967.
 14. Giotti, A.; Luzzi, S.; Spagnesi, S.; Zilletti, L. GABA_A and GABA_B receptor mediated effects in guinea-pig ileum. *Br. J. Pharmacol.* 78:469–478; 1983.
 15. Gower, A. J. Effects of acetylcholine agonists and antagonists on yawning and analgesia in the rat. *Eur. J. Pharmacol.* 139:79–89; 1987.
 16. Guyenet, P. G.; Agid, Y.; Javoy, F.; Beaujouan, J. C.; Rossier, J.; Glowinski, J. Effects of dopaminergic receptor agonists and antagonists on the activity of the neo-striatal cholinergic system. *Brain Res.* 84:227–244; 1974.
 17. Hadidian, J. Yawning in an old world monkey *Macaca Nigra* (Primates cercopithecidae). *Behavior* 75:133–147; 1980.
 18. Heusner, P. A. Yawning and associated phenomena. *Physiol. Rev.* 26:156–168; 1946.
 19. Holmgren, B.; Urbá-Holmgren, R. Yawning: A central cholinergic response under monoaminergic modulation. In: Tucek, S., ed. The cholinergic synapse. Amsterdam, The Netherlands: Elsevier Science Publishers; 1979:482. (Prog. Brain Res., vol. 49.)
 20. Holmgren, B.; Urbá-Holmgren, R. Interaction of cholinergic and dopaminergic influences on yawning behavior. *Acta Neurobiol. Exp.* 40:633–642; 1980.
 21. Jakobartl, L.; Huston, J. Spreading depression in hippocampus and neocortex of rats induced by ACTH 1–24. *Neurosci. Lett.* 5:189–192; 1977.
 22. Karbon, E. W.; Duman, R.; Enna, S. J. Biochemical identification of multiple GABA_B binding sites: association with noradrenergic terminals in rat forebrain. *Brain Res.* 274:393–396; 1983.
 23. Krogsgaard-Larsen, P.; Jacobsen, P.; Falch, E. Structure-activity requirements of the GABA receptor. In: Enna, S. J., ed. GABA receptors. Clifton, NJ: The Humana Press; 1983:149–176.
 24. Laping, N. J.; Ramirez, V. D. Prolactin induces yawning and the stretch-yawning syndrome in young adult male rats. *Horm. Behav.* 20:49–59; 1986.
 25. Laping, N. J.; Ramirez, V. D. Prolactin-induced yawning behavior requires an intact nigro-striatal dopamine system. *Pharmacol. Biochem. Behav.* 29:59–62; 1988.
 26. McGeer, E. G.; Staines, W. A.; McGeer, P. L. Neurotransmitters in the basal ganglia. *Can. J. Neurol. Sci.* 11:89–99; 1984.
 27. Marini, J. L. Serotonergic and dopaminergic effects on yawning in the cat. *Pharmacol. Biochem. Behav.* 15:711–715; 1981.
 28. Melis, M. R.; Argiolas, A.; Gessa, G. L. Oxytocin-induced penile erection and yawning: site of action in the brain. *Brain Res.* 398:259–265; 1986.
 29. Mogilnicka, E.; Klimek, V. Drugs affecting dopamine neurons and yawning behavior. *Pharmacol. Biochem. Behav.* 7:303–305; 1977.
 30. Park, M. R.; Lighthall, J. W.; Kitai, S. T. Recurrent inhibition in the rat neostriatum. *Brain Res.* 303:385–390; 1980.
 31. Scheel-Kruger, J. Dopamine-GABA interactions: evidence that GABA transmits, modulates and mediates dopaminergic functions in the basal ganglia and the limbic system. *Acta Neurol Scand. [Suppl.]* 107:9–50; 1986.
 32. Stahl, L.; Ungerstedt, U. Assessment of dopamine autoreceptor agonist properties of apomorphine, (+)-3-PPP and (–)-3-PPP by recording of yawning behavior in rat. *Eur. J. Pharmacol.* 98:307–310; 1984.
 33. Stoessl, A. J.; Dourish, C. T.; Iversen, S. D. Apomorphine-induced yawning in rats is abolished by bilateral 6-hydroxydopamine lesions of the substantia nigra. *Psychopharmacology (Berlin)* 93:336–342; 1987.
 34. Storm-Mathisen, J. Localization of transmitter candidates in the brain: the hippocampal formation as a model. *Prog. Neurobiol.* 8:119–181; 1977.
 35. Urbá-Holmgren, R.; Gonzalez, R. M.; Holmgren, B. Is yawning a cholinergic response? *Nature* 267:261–262; 1977.
 36. Urbá-Holmgren, R.; Holmgren, B.; Rodriguez, R.; Gonzalez, R. M. Serotonergic modulation of yawning. *Pharmacol. Biochem. Behav.* 11:371–372; 1979.
 37. Urbá-Holmgren, R.; Holmgren, B.; Anias, J. Pre and postsynaptic dopaminergic receptors involved in apomorphine-induced yawning. *Acta Neurobiol. Exp.* 42:115–125; 1982.
 38. Urbá-Holmgren, R.; Holmgren, B. Effects of metoclopramide on spontaneous and pharmacologically-induced yawning in the rat. *Bol. Estud. Med. Biol.* 35(3–4): 203–206; 1987.
 39. Ushijima, I.; Yamada, K.; Inoue, T.; Furukawa, T.; Noda, Y. Muscarinic and nicotinic effects on yawning and tongue protruding in the rat. *Pharmacol. Biochem. Behav.* 21:297–300; 1984.
 40. Ushijima, I.; Mizuki, Y.; Imaizumi, J.; Yamada, M.; Noda, Y.; Yamada, K.; Furukawa, T. Characteristics of yawning behavior induced by apomorphine, physostigmine and pilocarpine. *Arch. Int. Pharmacodyn. Ther.* 273(2):196–201; 1985.
 41. Wilson, C. J.; Groves, P. M. Spontaneous firing patterns of identified spiny neurons in the rat neostriatum. *Brain Res.* 220:67–80; 1981.
 42. Yamada, K.; Furukawa, T. Direct evidence for involvement of dopaminergic inhibition and cholinergic activation in yawning. *Psychopharmacology (Berlin)* 67:39–43; 1980.
 43. Yamada, K.; Tanaka, M.; Shibata, K.; Furukawa, T. Involvement of septal and striatal dopamine D-2 receptors in yawning behavior in rats. *Psychopharmacology (Berlin)* 90:9–13; 1986.